

GLOBAL EXISTENCE OF SMOOTH SOLUTIONS TO THREE-DIMENSIONAL TURBULENT FLOW EQUATIONS

DONGFEN BIAN AND BOLING GUO

ABSTRACT. In this paper we are concerned with the global existence of smooth solutions to the turbulent flow equations for compressible flows in $\mathbb{R}^3$. The global well-posedness is proved under the condition that the initial data are close to the standard equilibrium state in H^3 -framework. The proof relies on energy estimates about velocity, temperature, turbulent kinetic energy and rate of viscous dissipation. We use several new techniques to overcome the difficulties from the product of two functions and higher order norms. This is the first result concerning k - ε model equations.

AMS Subject Classification 2000: 35Q35, 35A01, 76F02.

Key words and phrases: Turbulent flow equations, compressible flows, k - ε model, smooth solution, global existence.

1. INTRODUCTION

We consider in this work the turbulent flow equations for compressible flows on $\mathbb{R}^3$,

$$\begin{cases} \rho_t + \operatorname{div}(\rho u) = 0, \\ (\rho u)_t + \operatorname{div}(\rho u \otimes u) - \Delta u - \nabla \operatorname{div} u + \nabla p = -\frac{2}{3} \nabla(\rho k), \\ (\rho h)_t + \operatorname{div}(\rho u h) - \Delta h = \frac{Dp}{Dt} + S_k, \\ (\rho k)_t + \operatorname{div}(\rho u k) - \Delta k = G - \rho \varepsilon, \\ (\rho \varepsilon)_t + \operatorname{div}(\rho u \varepsilon) - \Delta \varepsilon = \frac{C_1 G \varepsilon}{k} - \frac{C_2 \rho \varepsilon^2}{k}, \\ (\rho, u, h, k, \varepsilon)(x, t)|_{t=0} = (\rho_0(x), u_0(x), h_0(x), k_0(x), \varepsilon_0(x)), \end{cases} \quad (1.1)$$

with $S_k = [\mu(\frac{\partial u^i}{\partial x_j} + \frac{\partial u^j}{\partial x_i}) - \frac{2}{3} \delta_{ij} \mu \frac{\partial u^k}{\partial x_k}] \frac{\partial u^i}{\partial x_j} + \frac{\mu_t}{\rho^2} \frac{\partial p}{\partial x_j} \frac{\partial \rho}{\partial x_j}$, $G = \frac{\partial u^i}{\partial x_j} [\mu_e (\frac{\partial u^i}{\partial x_j} + \frac{\partial u^j}{\partial x_i}) - \frac{2}{3} \delta_{ij} (\rho k + \mu_e \frac{\partial u^k}{\partial x_k})]$, where $\delta_{ij} = 0$ if $i \neq j$, $\delta_{ij} = 1$ if $i = j$, and μ , μ_t , C_1 , C_2 are four positive constants satisfying $\mu + \mu_t = \mu_e$.

Here ρ , u , h , k and ε denote the density, velocity, total enthalpy, turbulent kinetic energy and rate of viscous dissipation, respectively. The pressure p is a smooth function of ρ . In this paper, without loss of generality, we have renormalized some constants to be 1.

The system (1.1) is formed by combining effect of turbulence on time-averaged Navier-Stokes equations with the k - ε model equations. Turbulence stands out as a prototype of multi-scale phenomenon that occurs in nature. It involves wide ranges of spatial and temporal scales which makes it very difficult to study analytically and prohibitively expensive to simulate computationally. Up to now, there are not any general theory suitable for the turbulent flows. Many, if not most, flows of engineering significance are turbulent, so the turbulent flow regime is not just of theoretical interest. Hence, research about the above system (1.1) is initial and very important.

There are many results about the simple model, such as classical Navier-Stokes equations and incompressible Navier-Stokes equations with variable density.

For classical Navier-Stokes equations (let $\rho = C$, $k = \varepsilon = 0$), there are many authors doing research about them, but the uniqueness and regularity of weak solutions for general initial data are open. In particular, Fujita and Kato [16, 17] proved the local well-posedness of 3D Navier-Stokes equations for large initial data and the global well-posedness for small initial data in the homogeneous Sobolev space $\dot{H}^{\frac{1}{2}}$. If taking u_0 small with respect to the viscosity in the Lebesgue space $L^N(\mathbb{R}^N)$, the classical Navier-Stokes equations are globally well-posed (see [18, 22]). This result has been adapted to the case of bounded domains by Y. Giga and T. Miyakawa in [19] and to the case of exterior domains by Y. Giga and H. Sohr in [20], and H. Iwashita in [21]. In the half-space case, the well-posedness issue has been studied by H. Kozono in [23] (see also [6] for results related to critical Besov spaces with negative index of regularity). In the absence of heat conduction, it was proved by Z. Xin that any non-zero smooth solution with initial compactly supported density would blow up in finite time(see [30]). As a reasonable starting point, we will therefore restrict our work to solutions such that ρ remains positive.

For incompressible Navier-Stokes equations with variable density (let $k = \varepsilon = 0$), there are also many results. Specially, O. Ladyzhenskaya and V. Solonnikov [24] studied the existence of strong (smooth) solutions with positive density whereas

the theory of global weak solutions with finite energy has been performed in the book [25]. Abidi [1] and Danchin-Mucha [8]-[10] have obtained several results about incompressible Navier-Stokes equations with variable density in the critical spaces. For more about well-posedness in other critical spaces, readers can refer to [2, 3, 4, 5, 7, 11, 12, 13, 14, 15, 26].

For the system (1.1), there is not any result. This paper is devoted to the study of the global existence of smooth solutions for the system (1.1) under suitable assumptions. Our result is expressed in the following.

Theorem 1.1. *Assume that initial data are close enough to the constant state $(\bar{\rho}, 0, 0, \bar{k}, 0)$, i.e. there exists a constant δ_0 such that if*

$$\|(\rho_0 - \bar{\rho}, u_0, h_0, k_0 - \bar{k}, \varepsilon_0)\|_{H^3(\mathbb{R}^3)} \leq \delta_0, \quad (1.2)$$

then the system (1.1) admits a unique smooth solution $(\rho, u, h, k, \varepsilon)$ such that for any $t \in [0, \infty)$,

$$\begin{aligned} & \|(\rho - \bar{\rho}, u, h, k - \bar{k}, \varepsilon)\|_{H^3}^2 + \int_0^t \|\nabla \rho\|_{H^2}^2 + \|(\nabla u, \nabla h, \nabla k, \nabla \varepsilon)\|_{H^3}^2 ds \\ & \leq C \|(\rho_0 - \bar{\rho}, u_0, h_0, k_0 - \bar{k}, \varepsilon_0)\|_{H^3}^2, \end{aligned}$$

where C is a positive constant.

Remark 1.2. The existence of the local solution for (1.1) can be obtained from the standard method based on the Banach theorem and contractivity of the operator defined by the linearization of the problem on a small time interval (see also [27, 28, 29]). Hence, we omit the local existence part for simplicity. The global existence of smooth solutions will be proved by extending the local solutions with respect to time based on a-priori global estimates.

Remark 1.3. We use several new techniques to overcome the difficulties from the product of two functions and higher order norms. In a separate paper we consider the convergence rate of smooth solution to the constant state.

Notation: Throughout the paper, C stands for a general constant, and may change from line to line. The norm $\|(A, B)\|_X$ is equivalent to $\|A\|_X + \|B\|_X$. The

notation $L^p(\mathbb{R}^3)$, $1 \leq p \leq \infty$, stands for the usual Lebesgue spaces on $\mathbb{R}^3$ and $\|\cdot\|_p$ denotes its L^p norm.

2. A PRIORI ESTIMATES

In this section we establish a-priori estimates of the solutions. we assume that $(v, u, \theta, k, \varepsilon)$ is a smooth solution to (1.1) on the time interval $(0, T)$ with $T > 0$. We shall establish the following proposition.

Proposition 2.1. *There exists a constant $\delta \ll 1$ such that if*

$$\sup_{0 \leq t \leq T} \|(\rho - \bar{\rho}, u, h, k - \bar{k}, \varepsilon)\|_{H^3} \leq \delta, \quad (2.1)$$

then for any $t \in [0, T]$, there exists a constant $C_1 > 1$ such that it holds

$$\begin{aligned} & \|(\rho - \bar{\rho}, u, h, k - \bar{k}, \varepsilon)\|_{H^3}^2 + \int_0^t \|\nabla \rho\|_{H^2}^2 + \|(\nabla u, \nabla h, \nabla k, \nabla \varepsilon)\|_{H^3}^2 ds \\ & \leq C_1 \|(\rho_0 - \bar{\rho}, u_0, h_0, k_0 - \bar{k}, \varepsilon_0)\|_{H^3}^2. \end{aligned} \quad (2.2)$$

Proof. First, let $\rho = a + \bar{\rho}$, $k = m + \bar{k}$, $f'(\rho) = \frac{p'(\rho)}{\rho}$, we rewrite the system (1.1) as follows:

$$\begin{cases} a_t + \operatorname{div}((a + \bar{\rho})u) = 0, \\ u_t + u \cdot \nabla u - \frac{1}{a + \bar{\rho}}(\Delta u + \nabla \operatorname{div} u) + \nabla[f(a + \bar{\rho}) - f(\bar{\rho})] \\ = -\frac{2}{3(a + \bar{\rho})}\nabla((a + \bar{\rho})(m + \bar{k})), \\ h_t + u \cdot \nabla h - \frac{1}{a + \bar{\rho}}\Delta h = -f'(a + \bar{\rho})(a + \bar{\rho})\operatorname{div} u + \frac{1}{a + \bar{\rho}}S_k, \\ m_t + u \cdot \nabla m - \frac{1}{a + \bar{\rho}}\Delta m = \frac{1}{a + \bar{\rho}}G - \varepsilon, \\ \varepsilon_t + u \cdot \nabla \varepsilon - \frac{1}{a + \bar{\rho}}\Delta \varepsilon = \frac{C_1 G \varepsilon}{(a + \bar{\rho})(m + \bar{k})} - \frac{C_2 \varepsilon^2}{m + \bar{k}}, \\ (a, u, h, m, \varepsilon)(x, t)|_{t=0} = (a_0(x), u_0(x), h_0(x), m_0(x), \varepsilon_0(x)), \end{cases} \quad (2.3)$$

with $S_k = [\mu(\frac{\partial u^i}{\partial x_j} + \frac{\partial u^j}{\partial x_i}) - \frac{2}{3}\delta_{ij}\mu\frac{\partial u^k}{\partial x_k}]\frac{\partial u^i}{\partial x_j} + \frac{\mu_t}{(a + \bar{\rho})^2}\frac{\partial p}{\partial x_j}\frac{\partial a}{\partial x_j}$, $G = \frac{\partial u^i}{\partial x_j}[\mu_e(\frac{\partial u^i}{\partial x_j} + \frac{\partial u^j}{\partial x_i}) - \frac{2}{3}\delta_{ij}((a + \bar{\rho})(m + \bar{k}) + \mu_e\frac{\partial u^k}{\partial x_k})]$.

From a priori assumption (2.1) and the Sobolev inequality together with the first equation of (2.3), we have

$$\sup_{x \in \mathbb{R}} |(a, a_t, \nabla a, u, \nabla u, h, \nabla h, m, \nabla m, \varepsilon, \nabla \varepsilon)| \leq C\|(a, u, h, m, \varepsilon)\|_{H^3} \leq C\delta. \quad (2.4)$$

Moreover,

$$\frac{\bar{\rho}}{2} \leq \rho = a + \bar{\rho} \leq 2\bar{\rho}, \quad \frac{\bar{k}}{2} \leq k = m + \bar{k} \leq 2\bar{k}, \quad (2.5)$$

and

$$0 < \frac{1}{C_0} \leq f'(\rho) \leq C_0 < \infty, \quad |f^{(n)}(\rho)| \leq C_0 \quad \text{for any positive integer } n, \quad (2.6)$$

with C_0 a positive constant.

In what follows, we will always use the smallness assumption of δ and (2.4)-(2.6).

We divide the a priori estimates into three steps.

Step 1: L^2 -norms of u , h , m , ε

Multiplying the second equation of (2.3) by u and integrating over $\mathbb{R}^3$, one can deduce that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|u\|_2^2 + \frac{1}{a + \bar{\rho}} (\|\nabla u\|_2^2 + \|\operatorname{div} u\|_2^2) \\ &= - \int_{\mathbb{R}^3} u \cdot \nabla u \cdot u dx + \int_{\mathbb{R}^3} (f(a + \bar{\rho}) - f(\bar{\rho})) \operatorname{div} u dx \\ &+ \int_{\mathbb{R}^3} \frac{1}{(a + \bar{\rho})^2} [(\nabla a \otimes u) : \nabla u + \operatorname{div} u (\nabla a \cdot u)] dx \\ &- \int_{\mathbb{R}^3} \frac{2}{3(a + \bar{\rho})} \nabla((a + \bar{\rho})(m + \bar{k})) \cdot u dx. \end{aligned} \quad (2.7)$$

With the help of Hölder's inequality, $\dot{W}^{1,2}(\mathbb{R}^3) \hookrightarrow L^6(\mathbb{R}^3)$ and (2.4), we estimate the right-hand side of (2.7) as

$$- \int_{\mathbb{R}^3} u \cdot \nabla u \cdot u dx \leq C \|u\|_6 \|\nabla u\|_2 \|u\|_3 \leq C \delta \|\nabla u\|_2^2, \quad (2.8)$$

$$\begin{aligned} & \int_{\mathbb{R}^3} \frac{1}{(a + \bar{\rho})^2} [(\nabla a \otimes u) : \nabla u + \operatorname{div} u (\nabla a \cdot u)] dx \\ & \leq C \|\nabla a\|_2 \|\nabla u\|_2 \|u\|_{L^\infty} \leq C \delta \|(\nabla a, \nabla u)\|_2^2, \end{aligned} \quad (2.9)$$

$$\begin{aligned} & - \int_{\mathbb{R}^3} \frac{2}{3(a + \bar{\rho})} \nabla((a + \bar{\rho})(m + \bar{k})) \cdot u dx \\ &= - \int_{\mathbb{R}^3} \frac{2(m + \bar{k})}{3(a + \bar{\rho})} \nabla a \cdot u - \frac{2}{3(a + \bar{\rho})} (a + \bar{\rho}) \nabla m \cdot u dx \\ & \leq C \delta \|(\nabla a, \nabla m)\|_2^2, \end{aligned} \quad (2.10)$$

$$\begin{aligned} & \int_{\mathbb{R}^3} (f(a + \bar{\rho}) - f(\bar{\rho})) \operatorname{div} u dx \\ &= - \int_{\mathbb{R}^3} [f(a + \bar{\rho}) - f(\bar{\rho})] \left(\frac{a_t + u \cdot \nabla a}{a + \bar{\rho}} \right) dx \\ &= - \frac{d}{dt} \int_{\mathbb{R}^3} F(a) dx - \int_{\mathbb{R}^3} \frac{f'(\bar{\rho} + \theta a)}{a + \bar{\rho}} a u \cdot \nabla a dx \\ & \leq - \frac{d}{dt} \int_{\mathbb{R}^3} F(a) dx + C \delta \|\nabla a\|_2^2, \end{aligned} \quad (2.11)$$

with $\theta \in (0, 1)$ and $F(a)$ is defined as

$$F(a) = \int_0^a \frac{f(s + \bar{\rho}) - f(\bar{\rho})}{s + \bar{\rho}} ds. \quad (2.12)$$

Combining (2.8)-(2.11) with (2.7), we get

$$\frac{1}{2} \frac{d}{dt} [\|u\|_2^2 + \int_{\mathbb{R}^3} F(a) dx] + \frac{1}{a + \bar{\rho}} (\|\nabla u\|_2^2 + \|\operatorname{div} u\|_2^2) \leq C\delta \|(\nabla a, \nabla u, \nabla m)\|_2^2. \quad (2.13)$$

Multiplying the energy equation, governing equation for turbulent kinetic energy k and ε -equation of (2.3) by h , m and ε respectively, and integrating them over the whole space $\mathbb{R}^3$, similarly we can get that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|h\|_2^2 + \frac{1}{a + \bar{\rho}} \|\nabla h\|_2^2 \\ &= \int_{\mathbb{R}^3} \frac{1}{(a + \bar{\rho})^2} \nabla a \cdot \nabla h \cdot h dx - \int_{\mathbb{R}^3} f'(a + \bar{\rho})(a + \bar{\rho}) \operatorname{div} u h dx \\ &+ \int_{\mathbb{R}^3} \frac{1}{a + \bar{\rho}} S_k \cdot h dx - \int_{\mathbb{R}^3} u \cdot \nabla h \cdot h dx, \end{aligned} \quad (2.14)$$

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|m\|_2^2 + \frac{1}{a + \bar{\rho}} \|\nabla m\|_2^2 \\ &= \int_{\mathbb{R}^3} \frac{1}{(a + \bar{\rho})^2} \nabla m \cdot \nabla a \cdot m dx - \int_{\mathbb{R}^3} \varepsilon \cdot m dx + \int_{\mathbb{R}^3} \frac{1}{a + \bar{\rho}} G \cdot m dx \\ &- \int_{\mathbb{R}^3} u \cdot \nabla m \cdot m dx, \end{aligned} \quad (2.15)$$

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|\varepsilon\|_2^2 + \frac{1}{a + \bar{\rho}} \|\nabla \varepsilon\|_2^2 \\ &= \int_{\mathbb{R}^3} \frac{1}{(a + \bar{\rho})^2} \nabla \varepsilon \cdot \nabla a \cdot \varepsilon dx + \int_{\mathbb{R}^3} \frac{C_1 G \varepsilon^2}{(a + \bar{\rho})(m + k)} dx \\ &- \int_{\mathbb{R}^3} \frac{C_2 \varepsilon^3}{m + k} dx - \int_{\mathbb{R}^3} u \cdot \nabla \varepsilon \cdot \varepsilon dx. \end{aligned} \quad (2.16)$$

A direct computation gives that

$$\int_{\mathbb{R}^3} \frac{1}{(a + \bar{\rho})^2} \nabla a \cdot \nabla h \cdot h dx \leq C \|\nabla a\|_2 \|\nabla h\|_2 \|h\|_{L^\infty} \leq C\delta \|(\nabla a, \nabla h)\|_2^2, \quad (2.17)$$

$$- \int_{\mathbb{R}^3} f'(a + \bar{\rho})(a + \bar{\rho}) \operatorname{div} u h dx \leq C \|\nabla a\|_2 \|\nabla u\|_2 \|h\|_3 \leq C\delta \|(\nabla a, \nabla u)\|_2^2, \quad (2.18)$$

$$\begin{aligned} & \int_{\mathbb{R}^3} \frac{1}{a + \bar{\rho}} S_k \cdot h dx \\ &= \int_{\mathbb{R}^3} \frac{1}{a + \bar{\rho}} \left\{ \left[\mu \left(\frac{\partial u^i}{\partial x_j} + \frac{\partial u^j}{\partial x_i} \right) - \frac{2}{3} \delta_{ij} \mu \frac{\partial u^k}{\partial x_k} \right] \frac{\partial u^i}{\partial x_j} + \frac{\mu_t}{(a + \bar{\rho})^2} \frac{\partial p}{\partial x_j} \frac{\partial a}{\partial x_j} \right\} \cdot h dx \\ &\leq C\delta \|(\nabla a, \nabla u)\|_2^2, \end{aligned} \quad (2.19)$$

$$- \int_{\mathbb{R}^3} u \cdot \nabla h \cdot h dx \leq C \|u\|_3 \|\nabla h\|_2 \|h\|_6 \leq C\delta \|\nabla h\|_2^2, \quad (2.20)$$

$$\int_{\mathbb{R}^3} \frac{1}{(a + \bar{\rho})^2} \nabla m \cdot \nabla a \cdot m dx \leq C \|\nabla m\|_2 \|\nabla a\|_2 \|m\|_{L^\infty} \leq C \delta \|(\nabla a, \nabla m)\|_2^2, \quad (2.21)$$

$$- \int_{\mathbb{R}^3} \varepsilon \cdot m dx \leq C \|\varepsilon\|_2 \|m\|_6 \|m + \bar{k}\|_6 \|m + \bar{k}\|_6 \leq C \delta \|\nabla m\|_2^2, \quad (2.22)$$

$$\begin{aligned} & \int_{\mathbb{R}^3} \frac{1}{a + \bar{\rho}} G \cdot m dx \\ &= \int_{\mathbb{R}^3} \frac{1}{a + \bar{\rho}} \frac{\partial u^i}{\partial x_j} [\mu_e (\frac{\partial u^i}{\partial x_j} + \frac{\partial u^j}{\partial x_i}) - \frac{2}{3} \delta_{ij} ((a + \bar{\rho})(m + \bar{k}) + \mu_e \frac{\partial u^k}{\partial x_k})] \cdot m dx \\ &\leq C \delta \|(\nabla u, \nabla m)\|_2^2, \end{aligned} \quad (2.23)$$

$$- \int_{\mathbb{R}^3} u \cdot \nabla m \cdot m dx \leq C \|u\|_3 \|\nabla m\|_2 \|m\|_6 \leq C \delta \|\nabla m\|_2^2, \quad (2.24)$$

$$\int_{\mathbb{R}^3} \frac{1}{(a + \bar{\rho})^2} \nabla \varepsilon \cdot \nabla a \cdot \varepsilon dx \leq C \|\nabla \varepsilon\|_2 \|\nabla a\|_2 \|\varepsilon\|_{L^\infty} \leq C \delta \|(\nabla a, \nabla \varepsilon)\|_2^2, \quad (2.25)$$

$$\begin{aligned} & \int_{\mathbb{R}^3} \frac{C_1 G \varepsilon^2}{(a + \bar{\rho})(m + \bar{k})} dx \\ &= \int_{\mathbb{R}^3} \frac{C_1 \varepsilon^2}{(a + \bar{\rho})(m + \bar{k})} \frac{\partial u^i}{\partial x_j} [\mu_e (\frac{\partial u^i}{\partial x_j} + \frac{\partial u^j}{\partial x_i}) - \frac{2}{3} \delta_{ij} ((a + \bar{\rho})(m + \bar{k}) \\ &+ \mu_e \frac{\partial u^k}{\partial x_k})] dx \leq C \delta \|(\nabla u, \nabla \varepsilon)\|_2^2, \end{aligned} \quad (2.26)$$

$$- \int_{\mathbb{R}^3} \frac{C_2 \varepsilon^3}{m + \bar{k}} dx \leq C \|\varepsilon\|_2 \|\varepsilon\|_6 \|\varepsilon\|_6 \|m + \bar{k}\|_6 \leq C \delta \|\nabla \varepsilon\|_2^2, \quad (2.27)$$

$$- \int_{\mathbb{R}^3} u \cdot \nabla \varepsilon \cdot \varepsilon dx \leq C \|u\|_3 \|\nabla \varepsilon\|_2 \|\varepsilon\|_6 \leq C \delta \|\nabla \varepsilon\|_2^2. \quad (2.28)$$

The estimates (2.17)-(2.28) together with (2.14)-(2.16) imply

$$\frac{1}{2} \frac{d}{dt} \|h\|_2^2 + \frac{1}{a + \bar{\rho}} \|\nabla h\|_2^2 \leq C \delta \|(\nabla a, \nabla u)\|_2^2, \quad (2.29)$$

$$\frac{1}{2} \frac{d}{dt} \|m\|_2^2 + \frac{1}{a + \bar{\rho}} \|\nabla m\|_2^2 \leq C \delta \|(\nabla a, \nabla u)\|_2^2, \quad (2.30)$$

$$\frac{1}{2} \frac{d}{dt} \|\varepsilon\|_2^2 + \frac{1}{a + \bar{\rho}} \|\nabla \varepsilon\|_2^2 \leq C \delta \|(\nabla a, \nabla u)\|_2^2. \quad (2.31)$$

Step 2: L^2 -norms of $\nabla^3 u$, $\nabla^3 h$, $\nabla^3 m$, $\nabla^3 \varepsilon$

Applying the differential operator ∂_{lmn} to the momentum equation of (2.3), then

multiplying it by $\partial_{lmn} u$, and integrating over $\mathbb{R}^3$, one gets

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|\partial_{lmn} u\|_2^2 \\ &= \int_{\mathbb{R}^3} \partial_{lmn} \left(\frac{1}{a + \bar{\rho}} \Delta u \right) \partial_{lmn} u dx + \int_{\mathbb{R}^3} \partial_{lmn} \left(\frac{1}{a + \bar{\rho}} \nabla \operatorname{div} u \right) \partial_{lmn} u dx \\ &- \int_{\mathbb{R}^3} \partial_{lmn} (u \cdot \nabla u) \cdot \partial_{lmn} u dx + \int_{\mathbb{R}^3} \partial_{lmn} [f(a + \bar{\rho}) - f(\bar{\rho})] \cdot \partial_{lmn} \operatorname{div} u dx \\ &- \frac{2}{3} \int_{\mathbb{R}^3} \partial_{lmn} \left[\frac{1}{a + \bar{\rho}} \nabla ((a + \bar{\rho})(m + \bar{k})) \right] \cdot \partial_{lmn} u dx. \end{aligned} \quad (2.32)$$

The first term on the right-hand side of (2.32) can be estimated as

$$\begin{aligned}
& \int_{\mathbb{R}^3} \partial_{lmn} \left(\frac{1}{a + \bar{\rho}} \Delta u \right) \partial_{lmn} u dx \\
&= - \int_{\mathbb{R}^3} \frac{1}{a + \bar{\rho}} |\partial_{lmn} \nabla u|^2 dx + \int_{\mathbb{R}^3} \frac{1}{(a + \bar{\rho})^2} \nabla \partial_{lmn} u : (\nabla a \otimes \partial_{lmn} u) dx \\
&+ \int_{\mathbb{R}^3} \left[- \frac{6}{(a + \bar{\rho})^4} \partial_l a \partial_m a \partial_n a \Delta u + \frac{2}{(a + \bar{\rho})^3} (\partial_{lm} a \partial_n a \Delta u + \partial_{ln} a \partial_m a \Delta u \right. \\
&+ \partial_{mn} a \partial_l a \Delta u + \partial_l a \partial_m a \partial_n \Delta u + \partial_l a \partial_n a \partial_m \Delta u + \partial_n a \partial_m a \partial_l \Delta u) \\
&- \frac{1}{(a + \bar{\rho})^2} (\partial_{lmn} a \Delta u + \partial_{lm} a \partial_n \Delta u + \partial_{ln} a \partial_m \Delta u + \partial_{mn} a \partial_l \Delta u \\
&+ \partial_l a \partial_{mn} \Delta u + \partial_m a \partial_{ln} \Delta u + \partial_n a \partial_{lm} \Delta u) \left. \right] \cdot \partial_{lmn} u dx \\
&\leq C\delta \|(\nabla^2 u, \nabla^3 u, \nabla^4 u, \nabla^2 a, \nabla^3 a)\|_2^2 - \int_{\mathbb{R}^3} \frac{1}{a + \bar{\rho}} |\partial_{lmn} \nabla u|^2 dx.
\end{aligned} \tag{2.33}$$

Similarly, we can estimate the second and third term on the right-hand side of (2.32) as

$$\begin{aligned}
& \int_{\mathbb{R}^3} \partial_{lmn} \left(\frac{1}{a + \bar{\rho}} \nabla \operatorname{div} u \right) \partial_{lmn} u dx \\
&\leq C\delta \|(\nabla^2 u, \nabla^3 u, \nabla^4 u, \nabla^2 a, \nabla^3 a)\|_2^2 - \int_{\mathbb{R}^3} \frac{1}{a + \bar{\rho}} |\partial_{lmn} \operatorname{div} u|^2 dx,
\end{aligned} \tag{2.34}$$

$$- \int_{\mathbb{R}^3} \partial_{lmn} (u \cdot \nabla u) \cdot \partial_{lmn} u dx \leq C\delta \|\nabla^3 u\|_2^2. \tag{2.35}$$

Now, let's estimate the fourth term on the right-hand side of (2.32) as

$$\begin{aligned}
& \int_{\mathbb{R}^3} \partial_{lmn} [f(a + \bar{\rho}) - f(\bar{\rho})] \cdot \partial_{lmn} \operatorname{div} u dx \\
&= - \int_{\mathbb{R}^3} \partial_{lmn} [f(a + \bar{\rho}) - f(\bar{\rho})] \partial_{lmn} \left(\frac{a_t + u \cdot \nabla a}{a + \bar{\rho}} \right) dx \\
&- \int_{\mathbb{R}^3} [f'''(\rho) \partial_l a \partial_m a \partial_n a + f''(\rho) \partial_{lm} a \partial_n a + f''(\rho) \partial_{ln} a \partial_m a \\
&+ f''(\rho) \partial_{mn} a \partial_l a + f'(\rho) \partial_{lmn} a] \times \left[-\frac{6}{(a + \bar{\rho})^4} \partial_l a \partial_m a \partial_n a (a_t + u \cdot \nabla a) \right. \\
&+ \frac{2}{(a + \bar{\rho})^3} (\partial_{lm} a \partial_n a a_t + \partial_{ln} a \partial_m a a_t + \partial_{mn} a \partial_l a a_t + \partial_l a \partial_m a \partial_n a_t \\
&+ \partial_l a \partial_n a \partial_m a_t + \partial_n a \partial_m a \partial_l a_t + u \cdot \nabla a \partial_{lm} a \partial_n a + u \cdot \nabla a \partial_{ln} a \partial_m a \\
&+ u \cdot \nabla a \partial_{mn} a \partial_l a + \partial_l u \cdot \nabla a \partial_m a \partial_n a + \partial_m u \cdot \nabla a \partial_l a \partial_n a + \partial_n u \cdot \nabla a \partial_l a \partial_m a \\
&+ u \cdot \partial_l \nabla a \partial_m a \partial_n a + u \cdot \partial_m \nabla a \partial_l a \partial_n a + u \cdot \partial_n \nabla a \partial_l a \partial_m a) \\
&- \frac{1}{(a + \bar{\rho})^2} (a_t \partial_{lmn} a + \partial_l a_t \partial_{mn} a + \partial_m a_t \partial_{ln} a + \partial_n a_t \partial_{lm} a + \partial_{lm} a_t \partial_n a \\
&+ \partial_{ln} a_t \partial_m a + \partial_{mn} a_t \partial_l a + u \cdot \nabla a \partial_{lmn} a + \partial_l u \cdot \nabla a \partial_{mn} a \\
&+ \partial_m u \cdot \nabla a \partial_{ln} a + \partial_n u \cdot \nabla a \partial_{lm} a + u \cdot \nabla \partial_l a \partial_{mn} a + u \cdot \nabla \partial_m a \partial_{ln} a \\
&+ u \cdot \nabla \partial_n a \partial_{lm} a + \partial_{lm} u \cdot \nabla a \partial_n a + \partial_{ln} u \cdot \nabla a \partial_m a + \partial_{mn} u \cdot \nabla a \partial_l a \\
&+ u \cdot \nabla \partial_{lm} a \partial_n a + u \cdot \nabla \partial_{ln} a \partial_m a + u \cdot \nabla \partial_{mn} a \partial_l a + \partial_l u \cdot \nabla \partial_m a \partial_n a \\
&+ \partial_l u \cdot \nabla \partial_n a \partial_m a + \partial_m u \cdot \nabla \partial_l a \partial_n a + \partial_m u \cdot \nabla \partial_n a \partial_l a + \partial_n u \cdot \nabla \partial_l a \partial_m a \\
&+ \partial_n u \cdot \nabla \partial_m a \partial_l a) + \frac{1}{a + \bar{\rho}} (\partial_{lmn} a_t + \partial_{lmn} u \cdot \nabla a + \partial_{lm} u \cdot \nabla \partial_n a \\
&+ \partial_{ln} u \cdot \nabla \partial_m a + \partial_{mn} u \cdot \nabla \partial_l a + \partial_l u \cdot \nabla \partial_{mn} a + \partial_m u \cdot \nabla \partial_{ln} a \\
&+ \partial_n u \cdot \nabla \partial_{lm} a + u \cdot \nabla \partial_{lmn} a)] dx \\
&\leq C \delta \|(\nabla a, \nabla^2 a, \nabla a_t, \nabla^3 a, \nabla^2 a_t, \nabla^2 u, \nabla^3 u)\|_2^2 - \int_{\mathbb{R}^3} [f'''(\rho) \partial_l a \partial_m a \partial_n a \\
&+ f''(\rho) \partial_{lm} a \partial_n a + f''(\rho) \partial_{ln} a \partial_m a + f''(\rho) \partial_{mn} a \partial_l a + f'(\rho) \partial_{lmn} a] \\
&\times \left[\frac{1}{a + \bar{\rho}} (\partial_{lmn} a_t + u \cdot \nabla \partial_{lmn} a) \right] dx.
\end{aligned} \tag{2.36}$$

Since

$$\begin{aligned}
- \int_{\mathbb{R}^3} \frac{1}{a + \bar{\rho}} f'''(\rho) \partial_l a \partial_m a \partial_n a \partial_{lmn} a_t dx &= \int_{\mathbb{R}^3} \partial_l \left(\frac{1}{a + \bar{\rho}} f'''(\rho) \partial_l a \partial_m a \partial_n a \right) \partial_{mn} a_t dx \\
&\leq C \delta \|(\nabla a, \nabla^2 a, \nabla^2 a_t)\|_2^2,
\end{aligned}$$

$$\begin{aligned}
- \int_{\mathbb{R}^3} \frac{1}{a + \bar{\rho}} f''(\rho) \partial_{lm} a \partial_n a \partial_{lmn} a_t dx &= \int_{\mathbb{R}^3} \partial_l \left(\frac{1}{a + \bar{\rho}} f''(\rho) \partial_{lm} a \partial_n a \right) \partial_{mn} a_t dx \\
&\leq C \delta \|(\nabla^2 a, \nabla^2 a_t, \nabla^3 a)\|_2^2, \\
- \int_{\mathbb{R}^3} \frac{1}{a + \bar{\rho}} f'''(\rho) \partial_l a \partial_m a \partial_n a u \cdot \nabla \partial_{lmn} a dx &= \int_{\mathbb{R}^3} \partial_j \left(\frac{1}{a + \bar{\rho}} f'''(\rho) \partial_l a \partial_m a \partial_n a u^j \right) \partial_{lmn} a dx \\
&\leq C \delta \|(\nabla^2 a, \nabla^3 a)\|_2^2,
\end{aligned}$$

and

$$\begin{aligned}
- \int_{\mathbb{R}^3} \frac{1}{a + \bar{\rho}} f''(\rho) \partial_{lm} a \partial_n a u \cdot \nabla \partial_{lmn} a dx &= \int_{\mathbb{R}^3} \partial_j \left(\frac{1}{a + \bar{\rho}} f''(\rho) \partial_{lm} a \partial_n a u^j \right) \partial_{lmn} a dx \\
&\leq C \delta \|\nabla^3 a\|_2^2,
\end{aligned}$$

we obtain

$$\begin{aligned}
&- \int_{\mathbb{R}^3} [f'''(\rho) \partial_l a \partial_m a \partial_n a + f''(\rho) \partial_{lm} a \partial_n a + f''(\rho) \partial_{ln} a \partial_m a + f''(\rho) \partial_{mn} a \partial_l a] \\
&\times \left[\frac{1}{a + \bar{\rho}} (\partial_{lmn} a_t + u \cdot \nabla \partial_{lmn} a) \right] \\
&\leq C \delta \|(\nabla a, \nabla^2 a, \nabla^2 a_t, \nabla^3 a)\|_2^2.
\end{aligned} \tag{2.37}$$

Finally,

$$\begin{aligned}
&- \int_{\mathbb{R}^3} \frac{f'(\rho)}{a + \bar{\rho}} \partial_{lmn} a (\partial_{lmn} a_t + u \cdot \nabla \partial_{lmn} a) dx \\
&= - \frac{1}{2} \int_{\mathbb{R}^3} \frac{f'(\rho)}{a + \bar{\rho}} (\partial_{lmn} a)_t^2 dx + \frac{1}{2} \int_{\mathbb{R}^3} \frac{f'(\rho)}{a + \bar{\rho}} (\partial_{lmn} a)^2 \operatorname{div} u + \frac{f''(\rho)}{a + \bar{\rho}} (\partial_{lmn} a)^2 u \cdot \nabla a \\
&- \frac{f'(\rho)}{(a + \bar{\rho})^2} (\partial_{lmn} a)^2 u \cdot \nabla a dx \\
&\leq - \frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^3} \frac{f'(\rho)}{a + \bar{\rho}} (\partial_{lmn} a)^2 dx + \frac{1}{2} \int_{\mathbb{R}^3} \left[\frac{f'(\rho)}{a + \bar{\rho}} \right]_t (\partial_{lmn} a)^2 dx + C \delta \|\nabla^3 a\|_2^2 \\
&\leq - \frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^3} \frac{f'(\rho)}{a + \bar{\rho}} (\partial_{lmn} a)^2 dx + C \delta \|\nabla^3 a\|_2^2,
\end{aligned}$$

together with (2.37), thus (2.36) can be replaced by

$$\begin{aligned}
&\int_{\mathbb{R}^3} \partial_{lmn} [f(a + \bar{\rho}) - f(\bar{\rho})] \cdot \partial_{lmn} \operatorname{div} u dx \\
&\leq C \delta \|(\nabla a, \nabla a_t, \nabla^2 a, \nabla^2 a_t, \nabla^3 a, \nabla^2 u, \nabla^3 u)\|_2^2 \\
&- \frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^3} \frac{f'(\rho)}{a + \bar{\rho}} (\partial_{lmn} a)^2 dx.
\end{aligned} \tag{2.38}$$

The last term on the right-hand side of (2.32) can be estimated as

$$\begin{aligned}
& -\frac{2}{3} \int_{\mathbb{R}^3} \partial_{lmn} \left[\frac{1}{a+\bar{\rho}} \nabla((a+\bar{\rho})(m+\bar{k})) \right] \cdot \partial_{lmn} u dx \\
& = \frac{2}{3} \int_{\mathbb{R}^3} \partial_{mn} \left[\frac{m+\bar{k}}{a+\bar{\rho}} \nabla a + \nabla m \right] \cdot \partial_{lmn} u dx \\
& = \frac{2}{3} \int_{\mathbb{R}^3} \left[\frac{2(m+\bar{k})}{(a+\bar{\rho})^3} \partial_m a \partial_n a \nabla a - \frac{1}{(a+\bar{\rho})^2} ((m+\bar{k}) \partial_{mn} a \nabla a + \partial_m a \partial_n m \nabla a \right. \\
& \quad + \partial_n a \partial_m m \nabla a + (m+\bar{k}) \partial_m a \partial_n \nabla a + (m+\bar{k}) \partial_n a \partial_m \nabla a) + \frac{1}{a+\bar{\rho}} (\partial_{mn} m \nabla a \\
& \quad + \partial_m m \nabla \partial_n a + \partial_n m \nabla \partial_m a + (m+\bar{k}) \nabla \partial_{mn} a) + \partial_{mn} \nabla m \left. \right] \cdot \partial_{lmn} u dx \\
& \leq C \delta \|(\nabla a, \nabla^2 a, \nabla^2 m, \nabla^3 m, \nabla^4 m, \nabla^4 u)\|_2^2 + C \int_{\mathbb{R}} (\nabla \partial_{mn} a)^2 dx,
\end{aligned}$$

which together with (2.33)-(2.35) and (2.38) gives

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} [\|\partial_{lmn} u\|_2^2 + \int_{\mathbb{R}^3} \frac{f'(\rho)}{a+\bar{\rho}} (\partial_{lmn} a)^2 dx] + \int_{\mathbb{R}^3} \frac{1}{a+\bar{\rho}} |\partial_{lmn} \nabla u|^2 dx \\
& + \int_{\mathbb{R}^3} \frac{1}{a+\bar{\rho}} |\partial_{lmn} \operatorname{div} u|^2 dx \\
& \leq C \delta \|(\nabla a, \nabla a_t, \nabla^2 a, \nabla^2 a_t, \nabla^3 a, \nabla^2 u, \nabla^3 u, \nabla^2 m, \nabla^3 m, \nabla^4 m)\|_2^2 \\
& + C \int_{\mathbb{R}} (\nabla \partial_{mn} a)^2 dx.
\end{aligned} \tag{2.39}$$

Applying ∂_{lmn} to the energy equation, k -equation and ε -equation of (2.3), multiplying the resulting equations by $\partial_{lmn} h$, $\partial_{lmn} m$ and $\partial_{lmn} \varepsilon$ respectively, then integrating them over the whole space $\mathbb{R}^3$, one can prove that

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \|\partial_{lmn} h\|_2^2 \\
& = \int_{\mathbb{R}^3} \partial_{lmn} \left(\frac{1}{a+\bar{\rho}} \Delta h \right) \cdot \partial_{lmn} h dx + \int_{\mathbb{R}^3} \partial_{lmn} \left(\frac{S_k}{a+\bar{\rho}} \right) \cdot \partial_{lmn} h dx \\
& - \int_{\mathbb{R}^3} \partial_{lmn} (f'(a+\bar{\rho})(a+\bar{\rho}) \operatorname{div} u) \cdot \partial_{lmn} h dx - \int_{\mathbb{R}^3} \partial_{lmn} (u \cdot \nabla h) \cdot \partial_{lmn} h dx,
\end{aligned} \tag{2.40}$$

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \|\partial_{lmn} m\|_2^2 \\
& = \int_{\mathbb{R}^3} \partial_{lmn} \left(\frac{1}{a+\bar{\rho}} \Delta m \right) \cdot \partial_{lmn} m dx - \int_{\mathbb{R}^3} \partial_{lmn} \varepsilon \cdot \partial_{lmn} m dx \\
& + \int_{\mathbb{R}^3} \partial_{lmn} \left(\frac{G}{a+\bar{\rho}} \right) \cdot \partial_{lmn} m dx - \int_{\mathbb{R}^3} \partial_{lmn} (u \cdot \nabla m) \cdot \partial_{lmn} m dx,
\end{aligned} \tag{2.41}$$

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \|\partial_{lmn} \varepsilon\|_2^2 \\
& = \int_{\mathbb{R}^3} \partial_{lmn} \left(\frac{1}{(a+\bar{\rho})^2} \Delta \varepsilon \right) \cdot \partial_{lmn} \varepsilon dx + \int_{\mathbb{R}^3} \partial_{lmn} \left(\frac{C_1 G \varepsilon}{(a+\bar{\rho})(m+\bar{k})} \right) \cdot \partial_{lmn} \varepsilon dx \\
& - \int_{\mathbb{R}^3} \partial_{lmn} \left(\frac{C_2 \varepsilon^2}{m+\bar{k}} \right) \cdot \partial_{lmn} \varepsilon dx - \int_{\mathbb{R}^3} \partial_{lmn} (u \cdot \nabla \varepsilon) \cdot \partial_{lmn} \varepsilon dx.
\end{aligned} \tag{2.42}$$

As same as the estimate (2.33), we can deduce

$$\begin{aligned} & \int_{\mathbb{R}^3} \partial_{lmn} \left(\frac{1}{a + \bar{\rho}} \Delta h \right) \partial_{lmn} h dx \\ & \leq C\delta \|(\nabla^2 h, \nabla^3 h, \nabla^4 h, \nabla^2 a, \nabla^3 a)\|_2^2 - \int_{\mathbb{R}^3} \frac{1}{a + \bar{\rho}} |\partial_{lmn} \nabla h|^2 dx, \end{aligned} \quad (2.43)$$

$$\begin{aligned} & \int_{\mathbb{R}^3} \partial_{lmn} \left(\frac{1}{a + \bar{\rho}} \Delta m \right) \partial_{lmn} m dx \\ & \leq C\delta \|(\nabla^2 m, \nabla^3 m, \nabla^4 m, \nabla^2 a, \nabla^3 a)\|_2^2 - \int_{\mathbb{R}^3} \frac{1}{a + \bar{\rho}} |\partial_{lmn} \nabla m|^2 dx, \end{aligned} \quad (2.44)$$

$$\begin{aligned} & \int_{\mathbb{R}^3} \partial_{lmn} \left(\frac{1}{a + \bar{\rho}} \Delta \varepsilon \right) \partial_{lmn} \varepsilon dx \\ & \leq C\delta \|(\nabla^2 \varepsilon, \nabla^3 \varepsilon, \nabla^4 \varepsilon, \nabla^2 a, \nabla^3 a)\|_2^2 - \int_{\mathbb{R}^3} \frac{1}{a + \bar{\rho}} |\partial_{lmn} \nabla \varepsilon|^2 dx. \end{aligned} \quad (2.45)$$

Again, from the Hölder inequality, (2.4)-(2.6), $\dot{W}^{1,2}(\mathbb{R}^3) \hookrightarrow L^6(\mathbb{R}^3)$ and $H^2(\mathbb{R}^3) \hookrightarrow L^\infty(\mathbb{R}^3)$, we get

$$\begin{aligned} & \int_{\mathbb{R}^3} \partial_{lmn} \left(\frac{S_k}{a + \bar{\rho}} \right) \cdot \partial_{lmn} h dx \\ & = \int_{\mathbb{R}^3} \partial_{lmn} \left\{ \frac{1}{a + \bar{\rho}} \left[\left(\mu \left(\frac{\partial u^i}{\partial x_j} + \frac{\partial u^j}{\partial x_i} \right) - \frac{2}{3} \delta_{ij} \mu \frac{\partial u^k}{\partial x_k} \right) \frac{\partial u^i}{\partial x_j} + \frac{\mu_t}{(a + \bar{\rho})^2} \frac{\partial p}{\partial x_j} \frac{\partial a}{\partial x_j} \right] \right\} \cdot \partial_{lmn} h dx \\ & \leq C\delta \|(\nabla^3 a, \nabla^3 u, \nabla^4 u, \nabla^3 h, \nabla^4 h)\|_2^2, \\ & \int_{\mathbb{R}^3} \partial_{lmn} \left(\frac{G}{a + \bar{\rho}} \right) \cdot \partial_{lmn} m dx \\ & = \int_{\mathbb{R}^3} \partial_{lmn} \left\{ \frac{1}{a + \bar{\rho}} \frac{\partial u^i}{\partial x_j} \left[\mu_e \left(\frac{\partial u^i}{\partial x_j} + \frac{\partial u^j}{\partial x_i} \right) - \frac{2}{3} \delta_{ij} ((a + \bar{\rho})(m + \bar{k}) + \mu_e \frac{\partial u^k}{\partial x_k}) \right] \right\} \cdot \partial_{lmn} m dx \\ & \leq C\delta \|(\nabla^2 a, \nabla^3 a, \nabla^3 m, \nabla^4 m, \nabla^3 u, \nabla^4 u)\|_2^2, \\ & \int_{\mathbb{R}^3} \partial_{lmn} \left(\frac{C_1 G \varepsilon}{(a + \bar{\rho})(m + \bar{k})} \right) \cdot \partial_{lmn} \varepsilon dx \\ & = \int_{\mathbb{R}^3} \partial_{lmn} \left\{ \frac{C_1 \varepsilon}{(a + \bar{\rho})(m + \bar{k})} \frac{\partial u^i}{\partial x_j} \left[\mu_e \left(\frac{\partial u^i}{\partial x_j} + \frac{\partial u^j}{\partial x_i} \right) - \frac{2}{3} \delta_{ij} ((a + \bar{\rho})(m + \bar{k}) \right. \right. \\ & \quad \left. \left. + \mu_e \frac{\partial u^k}{\partial x_k}) \right] \right\} \cdot \partial_{lmn} \varepsilon dx \\ & \leq C\delta \|(\nabla^3 a, \nabla^3 u, \nabla^4 u, \nabla^3 m, \nabla^3 \varepsilon)\|_2^2, \\ & - \int_{\mathbb{R}^3} \partial_{lmn} (f'(a + \bar{\rho})(a + \bar{\rho}) \operatorname{div} u) \cdot \partial_{lmn} h dx \leq C\delta \|(\nabla^3 a, \nabla^3 u, \nabla^4 u, \nabla^3 h, \nabla^4 h)\|_2^2, \\ & - \int_{\mathbb{R}^3} \partial_{lmn} (u \cdot \nabla h) \cdot \partial_{lmn} h dx \leq C\delta \|(\nabla^3 u, \nabla^3 h, \nabla^4 h)\|_2^2, \\ & - \int_{\mathbb{R}^3} \partial_{lmn} \varepsilon \cdot \partial_{lmn} m dx \leq C\delta \|(\nabla^3 \varepsilon, \nabla^3 m, \nabla^4 m)\|_2^2, \\ & - \int_{\mathbb{R}^3} \partial_{lmn} (u \cdot \nabla m) \cdot \partial_{lmn} m dx \leq C\delta \|(\nabla^3 m, \nabla^4 m, \nabla^3 u)\|_2^2, \end{aligned}$$

$$\begin{aligned}
& - \int_{\mathbb{R}^3} \partial_{lmn} \left(\frac{C_2 \varepsilon^2}{m+k} \right) \cdot \partial_{lmn} \varepsilon dx \leq C\delta \|(\nabla^3 m, \nabla^3 \varepsilon)\|_2^2, \\
& - \int_{\mathbb{R}^3} \partial_{lmn} (u \cdot \nabla \varepsilon) \cdot \partial_{lmn} \varepsilon dx \leq C \|(\nabla^3 \varepsilon, \nabla^4 \varepsilon, \nabla^3 u)\|_2^2.
\end{aligned}$$

Incorporating the above estimates and (2.40)-(2.42) yields that

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \|\partial_{lmn} h\|_2^2 + \int_{\mathbb{R}^3} \frac{1}{a+\bar{\rho}} |\partial_{lmn} \nabla h|^2 dx \\
& \leq C\delta \|(\nabla^2 a, \nabla^3 a, \nabla^2 h, \nabla^3 h, \nabla^3 u, \nabla^4 u)\|_2^2,
\end{aligned} \tag{2.46}$$

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \|\partial_{lmn} m\|_2^2 + \int_{\mathbb{R}^3} \frac{1}{a+\bar{\rho}} |\partial_{lmn} \nabla m|^2 dx \\
& \leq C\delta \|(\nabla^2 a, \nabla^3 a, \nabla^2 m, \nabla^3 m, \nabla^3 \varepsilon, \nabla^3 u, \nabla^4 u)\|_2^2,
\end{aligned} \tag{2.47}$$

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \|\partial_{lmn} \varepsilon\|_2^2 + \int_{\mathbb{R}^3} \frac{1}{a+\bar{\rho}} |\partial_{lmn} \nabla \varepsilon|^2 dx \\
& \leq C\delta \|(\nabla^2 a, \nabla^3 a, \nabla^3 m, \nabla^2 \varepsilon, \nabla^3 \varepsilon, \nabla^3 u, \nabla^4 u)\|_2^2.
\end{aligned} \tag{2.48}$$

Step 3: L^2 -norms of $\nabla a, \nabla^3 a$

We first estimate for ∇a . For this purpose, we calculate as

$$\begin{aligned}
& \int_{\mathbb{R}^3} \left[\frac{1}{2} |\nabla a|^2 + \frac{(a+\bar{\rho})^2}{2} \nabla a \cdot u \right]_t \\
& = \int_{\mathbb{R}^3} \nabla a \cdot \nabla a_t + (a+\bar{\rho}) a_t \nabla a \cdot u + \frac{(a+\bar{\rho})^2}{2} \nabla a_t \cdot u + \frac{(a+\bar{\rho})^2}{2} \nabla a \cdot u_t dx.
\end{aligned} \tag{2.49}$$

The first term of the right-hand side of (2.49) can be estimated as follows:

$$\begin{aligned}
& \int_{\mathbb{R}^3} \nabla a \cdot \nabla a_t dx = - \int_{\mathbb{R}^3} \nabla a \cdot \nabla \operatorname{div}[(a+\bar{\rho})u] dx \\
& = - \int_{\mathbb{R}^3} \nabla a \cdot (\nabla^2 a \cdot u) + \nabla a \cdot (\nabla a \cdot \nabla u) \\
& \quad + (a+\bar{\rho}) \nabla a \cdot \nabla \operatorname{div} u + |\nabla a|^2 \operatorname{div} u dx \\
& \leq C\delta \|\nabla a\|_2^2 - \int_{\mathbb{R}^3} (a+\bar{\rho}) \nabla a \cdot \nabla \operatorname{div} u dx.
\end{aligned} \tag{2.50}$$

Also, we estimate the rest three terms as

$$\int_{\mathbb{R}^3} (a+\bar{\rho}) a_t \nabla a \cdot u dx \leq C\delta \|(a_t, \nabla a)\|_2^2 \leq C\delta \|(\nabla a, \nabla u)\|_2^2, \tag{2.51}$$

$$\int_{\mathbb{R}^3} \frac{(a+\bar{\rho})^2}{2} \nabla a_t \cdot u dx \leq C\delta \|(\nabla a, \nabla u)\|_2^2, \tag{2.52}$$

$$\begin{aligned}
& \int_{\mathbb{R}^3} \frac{(a + \bar{\rho})^2}{2} \nabla a \cdot u_t dx \\
&= \int_{\mathbb{R}^3} \frac{(a + \bar{\rho})^2}{2} \nabla a \cdot \left(-u \cdot \nabla u + \frac{1}{a + \bar{\rho}} \Delta u + \frac{1}{a + \bar{\rho}} \nabla \operatorname{div} u \right. \\
&\quad \left. - \nabla [f(a + \bar{\rho}) - f(\bar{\rho})] - \frac{2}{3(a + \bar{\rho})} \nabla((a + \bar{\rho})(m + \bar{k})) \right) dx \\
&\leq C\delta \|(\nabla a, \nabla u)\|_2^2 - \int_{\mathbb{R}^3} \frac{(a + \bar{\rho})(m + \bar{k})}{3} |\nabla a|^2 dx \\
&\quad - \int_{\mathbb{R}^3} \frac{(a + \bar{\rho})^2}{2} f'(\rho) |\nabla a|^2 dx + \int_{\mathbb{R}^3} \frac{a + \bar{\rho}}{2} \nabla a \cdot (\Delta u + \nabla \operatorname{div} u) dx.
\end{aligned} \tag{2.53}$$

Since

$$\begin{aligned}
& \int_{\mathbb{R}^3} \frac{a + \bar{\rho}}{2} \nabla a \cdot (\Delta u - \nabla \operatorname{div} u) dx = \int_{\mathbb{R}^3} \frac{a + \bar{\rho}}{2} \partial_i a \cdot (\partial_{jj} u^i - \partial_i \partial_j u^j) dx \\
&= \int_{\mathbb{R}^3} -\frac{1}{2} \partial_i a \partial_j a \partial_j u^i - \frac{a + \bar{\rho}}{2} \partial_{ij} a \partial_j u^i + \frac{1}{2} \partial_i a \partial_j a \partial_i u^j + \frac{a + \bar{\rho}}{2} \partial_{ij} a \partial_i u^j dx = 0,
\end{aligned}$$

together with (2.49)-(2.53) and (2.5)-(2.6), we show that

$$\int_{\mathbb{R}^3} \left[\frac{1}{2} |\nabla a|^2 + \frac{(a + \bar{\rho})^2}{2} \nabla a \cdot u \right]_t + C \|\nabla a\|_2^2 \leq C\delta \|\nabla u\|_2^2. \tag{2.54}$$

Now, we turn to estimate for $\nabla^3 a$. Almost parallel to the inequality (2.49), we get

$$\begin{aligned}
& \int_{\mathbb{R}^3} \left[\frac{1}{2} |\partial_{lmn} a|^2 + \frac{(a + \bar{\rho})^2}{2} \partial_{lmn} a \partial_{lm} u^n \right]_t dx \\
&= \int_{\mathbb{R}^3} -\partial_{lmn} a \partial_{lmn} \operatorname{div}[(a + \bar{\rho})u] + (a + \bar{\rho}) a_t \partial_{lmn} a \partial_{lm} u^n \\
&\quad - \frac{(a + \bar{\rho})^2}{2} \partial_{lmn} \operatorname{div}[(a + \bar{\rho})u] \partial_{lm} u^n + \frac{(a + \bar{\rho})^2}{2} \partial_{lmn} a \partial_{lm} u_t^n dx.
\end{aligned} \tag{2.55}$$

We estimate the right-hand side of (2.55) as follows:

$$\begin{aligned}
& \int_{\mathbb{R}^3} -\partial_{lmn} a \partial_{lmn} \operatorname{div}[(a + \bar{\rho})u] dx \\
&= \int_{\mathbb{R}^3} -\partial_{lmn} a [u \cdot \nabla \partial_{lmn} a + \partial_l u \cdot \nabla \partial_{mn} a + \partial_m u \cdot \nabla \partial_{ln} a + \partial_n u \cdot \nabla \partial_{lm} a \\
&\quad + \partial_{lm} u \cdot \nabla \partial_n a + \partial_{ln} u \cdot \nabla \partial_m a + \partial_{mn} u \cdot \nabla \partial_l a + \partial_{lmn} u \cdot \nabla a + \partial_{lmn} a \operatorname{div} u \\
&\quad + \partial_{lm} a \partial_n \operatorname{div} u + \partial_{ln} a \partial_m \operatorname{div} u + \partial_{mn} a \partial_l \operatorname{div} u + \partial_l a \partial_{mn} \operatorname{div} u + \partial_m a \partial_{ln} \operatorname{div} u \\
&\quad + \partial_n a \partial_{lm} \operatorname{div} u + (a + \bar{\rho}) \partial_{lmn} \operatorname{div} u] dx \\
&\leq C\delta \|(\nabla^3 a, \nabla^3 u)\|_2^2 - \int_{\mathbb{R}^3} (a + \bar{\rho}) \partial_{lmn} a \partial_{lmn} \operatorname{div} u dx,
\end{aligned} \tag{2.56}$$

$$\int_{\mathbb{R}^3} (a + \bar{\rho}) a_t \partial_{lmn} a \partial_{lm} u^n dx \leq C\delta \|(\nabla^3 a, \nabla^2 u)\|_2^2, \tag{2.57}$$

$$\begin{aligned}
& - \int_{\mathbb{R}^3} \frac{(a + \bar{\rho})^2}{2} \partial_{lmn} \operatorname{div}[(a + \bar{\rho})u] \partial_{lm} u^n dx \\
& = - \int_{\mathbb{R}^3} \frac{(a + \bar{\rho})^2}{2} \partial_{lm} u^n [\partial_{lmn}(u \cdot \nabla a) + \partial_{lmn} a \operatorname{div} u + \partial_{lm} a \partial_n \operatorname{div} u \\
& \quad + \partial_{ln} a \partial_m \operatorname{div} u + \partial_{mn} a \partial_l \operatorname{div} u + \partial_l a \partial_{mn} \operatorname{div} u + \partial_m a \partial_{ln} \operatorname{div} u + \partial_n a \partial_{lm} \operatorname{div} u \\
& \quad + (a + \bar{\rho}) \partial_{lmn} \operatorname{div} u] dx \\
& \leq C\delta \|(\nabla^2 a, \nabla^3 a, \nabla^2 u, \nabla^3 u, \nabla^4 u)\|_2^2,
\end{aligned} \tag{2.58}$$

$$\begin{aligned}
& \int_{\mathbb{R}^3} \frac{(a + \bar{\rho})^2}{2} \partial_{lmn} a \partial_{lm} u_t^n dx \\
& = \int_{\mathbb{R}^3} \frac{(a + \bar{\rho})^2}{2} \partial_{lmn} a \partial_{lm} [-u^i \partial_i u^n + \frac{1}{a + \bar{\rho}} \partial_{ii} u^n + \frac{1}{a + \bar{\rho}} \partial_{in} u^i \\
& \quad - \partial_n(f(a + \bar{\rho}) - f(\bar{\rho})) - \frac{2}{3(a + \bar{\rho})} \partial_n((a + \bar{\rho})(m + \bar{k}))] dx \\
& \leq C\delta \|(\nabla^2 a, \nabla^3 a, \nabla^2 u, \nabla^3 u, \nabla^3 m, \nabla^4 m)\|_2^2 \\
& \quad + \int_{\mathbb{R}^3} \frac{a + \bar{\rho}}{2} \partial_{lmn} a (\partial_{lmii} u^n + \partial_{lmni} u^i) dx - \int_{\mathbb{R}^3} \frac{(a + \bar{\rho})^2}{2} f'(\rho) (\partial_{lmn} a)^2 dx \\
& \quad - \int_{\mathbb{R}^3} \frac{(a + \bar{\rho})(m + \bar{k})}{3} (\partial_{lmn} a)^2 dx.
\end{aligned} \tag{2.59}$$

Noting that

$$\begin{aligned}
& - \int_{\mathbb{R}} (a + \bar{\rho}) \partial_{lmn} a \partial_{lmn} \operatorname{div} u dx + \int_{\mathbb{R}^3} \frac{a + \bar{\rho}}{2} \partial_{lmn} a (\partial_{lmii} u^n + \partial_{lmni} u^i) dx \\
& = \int_{\mathbb{R}^3} \frac{a + \bar{\rho}}{2} \partial_{lmn} a (\partial_{lmii} u^n - \partial_{lmni} u^i) dx \\
& = -\frac{1}{2} \int_{\mathbb{R}^3} \partial_i a \partial_{lmn} a \partial_{lmi} u^n + (a + \bar{\rho}) \partial_{lmni} a \partial_{lmi} u^n - \partial_i a \partial_{lmn} a \partial_{lmn} u^i \\
& \quad - (a + \bar{\rho}) \partial_{lmni} a \partial_{lmn} u^i dx \\
& = -\frac{1}{2} \int_{\mathbb{R}^3} \partial_i a \partial_{lmn} a \partial_{lmi} u^n - \partial_i a \partial_{lmn} a \partial_{lmn} u^i dx \\
& \leq C\delta \|(\nabla^3 a, \nabla^3 u)\|_2^2,
\end{aligned}$$

based on the estimates (2.55)-(2.59), with the help of (2.5)-(2.6) and the interpolation inequality, one obtains

$$\begin{aligned}
& \int_{\mathbb{R}^3} [\frac{1}{2} |\partial_{lmn} a|^2 + \frac{(a + \bar{\rho})^2}{2} \partial_{lmn} a \partial_{lm} u^n]_t dx + C \|\nabla^3 a\|_2^2 \\
& \leq C\delta \|(\nabla^2 a, \nabla^2 u, \nabla^3 u, \nabla^4 u, \nabla^3 m, \nabla^4 m)\|_2^2.
\end{aligned} \tag{2.60}$$

Step 4: Conclusion

Consequently, multiplying (2.39) by a appropriate small constants α , together with (2.13), (2.29)-(2.31), (2.46)-(2.48), (2.54) and (2.60), we have

$$\begin{aligned}
& \frac{d}{dt} \{ \| (u, h, \nabla^3 h, m, \nabla^3 m, \varepsilon, \nabla^3 \varepsilon) \|_2^2 + \int_{\mathbb{R}^3} F(a) dx + \alpha [\| \nabla^3 u \|_2^2 \\
& + \int_{\mathbb{R}^3} \frac{f'(\rho)}{a + \bar{\rho}} (\partial_{lmn} a)^2 dx] + \int_{\mathbb{R}^3} [\frac{1}{2} |\nabla a|^2 + \frac{(a + \bar{\rho})^2}{2} \nabla a \cdot u \\
& + \frac{1}{2} |\partial_{lmn} a|^2 + \frac{(a + \bar{\rho})^2}{2} \partial_{lmn} a \partial_{lm} u^n] dx \} \\
& + C(\alpha) \| (\nabla a, \nabla^3 a, \nabla u, \nabla^4 u, \nabla h, \nabla^4 h, \nabla m, \nabla^4 m, \nabla \varepsilon, \nabla^4 \varepsilon) \|_2^2 \\
& \leq 0,
\end{aligned} \tag{2.61}$$

where we have used the fact that

$$|\nabla^i a_t| \leq C \sum_{k=1}^{i+1} (|\nabla^k a| + |\nabla^k u|), \quad i = 1, 2.$$

Integrating the inequality (2.61), from (2.4)-(2.6), (2.12) and the smallness of α , we can finish the proof of Proposition 2.1. $\square$

3. PROOF OF GLOBAL EXISTENCE

We will finish the proof of Theorem 1.1 in this section. First, let's state the local existence. Since it can be proved in a standard way as that in [27, 29], we omit the proof.

Proposition 3.1. *Under the assumption of the Theorem 1.1, then there exists a constant $T > 0$ such that the system (2.3) admits a unique smooth solution $(\rho, u, h, k, \varepsilon)$ which satisfies that there exists a constant $C_2 > 1$ such that for any $t \in [0, T]$,*

$$\begin{aligned}
& \| (\rho - \bar{\rho}, u, h, k - \bar{k}, \varepsilon) \|_{H^3}^2 + \int_0^t (\| \nabla \rho \|_{H^2}^2 + \| (\nabla u, \nabla h, \nabla k, \nabla \varepsilon) \|_{H^3}^2) ds \\
& \leq C_2 \| (\rho_0 - \bar{\rho}, u_0, h_0, k_0 - \bar{k}, \varepsilon_0) \|_{H^3}^2.
\end{aligned} \tag{3.1}$$

In the following, by a continued argument, combining the local existence and the a priori estimates proposition, we will prove the global existence of smooth solutions.

First, suppose

$$E_0 = \| (\rho_0 - \bar{\rho}, u_0, h_0, k_0 - \bar{k}, \varepsilon_0) \|_{H^3} < \min(\delta / \sqrt{C_2}, \delta / \sqrt{C_1 C_2}), \tag{3.2}$$

where δ is defined in Proposition 2.1. Since the initial data satisfy $E_0 < \delta/\sqrt{C_2}$, then by Proposition 3.1, there exists a constant $T^* > 0$ such that there exists a unique solution on $[0, T_*]$ satisfying

$$E_1 := \sup_{0 \leq t \leq T_1} \|(\rho - \bar{\rho}, u, h, k - \bar{k}, \varepsilon)\|_{H^3} \leq \sqrt{C_2} E_0. \quad (3.3)$$

Therefore, using the inequality $E_0 < \delta/\sqrt{C_1 C_2}$, from Proposition 2.1, we have

$$E_1 \leq \sqrt{C_1} E_0 < \delta/\sqrt{C_2}. \quad (3.4)$$

Notice that T^* depends only on E_0 . Starting from T^* , then the initial problem (2.3) with initial data $(\rho, u, h, k, \varepsilon)(T^*)$ still has a unique solution on $[T^*, 2T^*]$, and from Proposition 3.1, we get

$$\sup_{T^* \leq t \leq 2T^*} \|(\rho - \bar{\rho}, u, h, k - \bar{k}, \varepsilon)\|_{H^3} \leq \sqrt{C_2} E_1 \leq \sqrt{C_1 C_2} E_0 \leq \delta.$$

Again from Proposition 2.1, one can deduce

$$E_2 = \sup_{0 \leq t \leq 2T^*} \|(\rho - \bar{\rho}, u, h, k - \bar{k}, \varepsilon)\|_{H^3} \leq \sqrt{C_1} E_0 < \delta/\sqrt{C_2}.$$

Repeating the procedure for $0 \leq t \leq NT_*$, $N = 1, 2, 3, \dots$, we can extend the local solution to infinity as far as the initial data are small enough such that $E_0 \leq \min(\delta/\sqrt{C_2}, \delta/\sqrt{C_1 C_2})$. Thus the proof of Theorem 1.1 is complete. $\square$

REFERENCES

- [1] H. Abidi, Équation de Navier-Stokes avec densité et viscosité variables dans l'espace critique, *Rev. Mat. Iberoam.* 23(2)(2007) 537–586.
- [2] D. F. Bian, B. Q. Yuan, Local well-posedness in critical spaces for compressible MHD equations, (2010) 1–30, preprint.
- [3] D. F. Bian, B. Q. Yuan, Well-posedness in super critical Besov spaces for compressible MHD equations, *Int. J. Dynamical Systems and Differential Equations* 3(2011) 383–399.
- [4] M. Cannone, Ondelettes, Paraproducts et Navier-Stokes equation, *Nouveaux essais*, Diderot éditeurs, Paris, 1995.
- [5] M. Cannone, Harmonic analysis tools for solving the incompressible Navier-Stokes equations, *Handbook of Mathematical fluid Dynamics*, vol. III, North-Holland, Amsterdam, 2004.
- [6] M. Cannone, F. Planchon, M. Schonbek, Strong solutions to the incompressible Navier-Stokes equations in the half-space, *Comm. Partial Differential Equations* 25(5-6)(2000) 903–924.
- [7] Q. Chen, C. Miao, Z. Zhang, Well-posedness in critical spaces for compressible Navier-Stokes equations with density dependent viscosities, *Rev. Mat. Iberoamericana* 26(2010) 915–946.
- [8] R. Danchin, Density-dependent incompressible viscous fluids in critical spaces, *Proc. Roy. Soc. Edinburgh Sect. A* 133(2003) 1311–1334.
- [9] R. Danchin, P. B. Mucha, A critical functional framework for the inhomogeneous Navier-Stokes equations in the half-space, *J. Funct. Anal.* 256(3)(2009) 881–927.
- [10] R. Danchin and P. B. Mucha, Critical functional framework and maximal regularity in action on systems of incompressible flows, in progress.
- [11] R. Danchin, Local theory in critical spaces for compressible viscous and heat-conductive Gases, *Comm. Partial Differential Equations* 26(2001) 1183–1233.

- [12] R. Danchin, Global existence in critical spaces for flows of compressible viscous and heat-conductive Gases, Arch. Rational Mech. Anal. 160(2001) 1–39.
- [13] R. Danchin, Global existence in critical spaces for compressible Navier-Stokes equations, Invent. Math. Anal. 141(2000) 579–614.
- [14] R. Danchin, On the uniqueness in critical spaces for Compressible Navier-Stokes equations, Nonlinear Differential Equations Appl. 12(2005) 111–128.
- [15] R. Danchin, Well-posedness in critical spaces for barotropic viscous fluids with truly not constant density, Comm. Partial Differential Equations 32(2007) 1373–1397.
- [16] H. Fujita and T. Kato, On Navier-Stokes initial value problem, I. Arch. Rational Mech. Anal. 16(1964) 269–315.
- [17] H. Fujita, T. Kato, On the nonstationary Navier-Stokes system, Rend. Sem. Mat. Univ. Padova 32(1962) 243–260.
- [18] Y. Giga, Solutions for semilinear parabolic equations in L^p and regularity of weak solutions of the Navier-Stokes system, J. Differential Equations 62(2)(1986) 186–212.
- [19] Y. Giga, T. Miyakawa, Solutions in L^r of the Navier-Stokes initial value problem, Arch. Ration. Mech. Anal. 89(1985) 267–281.
- [20] Y. Giga, H. Sohr, Abstract L^p estimates for the cauchy problem with applications to the Navier-Stokes equations in exterior domains, J. Funct. Anal. 102(1991) 72–94.
- [21] H. Iwashita, L^q - L^r estimates for solutions of the nonstationary Stokes equations in an exterior domain and the Navier-Stokes initial value problems in L^q spaces, Math. Ann. 285(2)(1989) 265–288.
- [22] T. Kato, Strong L^p -solutions of Navier-Stokes equations in $\mathbb{R}^N$ with applications to weak solutions, Math. Z 187(1984) 471–480.
- [23] H. Kozono, Global L^N -solution and its decay property for the Navier-Stokes equations in half-space $\mathbb{R}_+^N$, J. Differential Equations 79(1)(1989) 79–88.
- [24] O. Ladyzhenskaya, V. Solonnikov, The unique solvability of an initial-boundary value problem for viscous incompressible inhomogeneous fluids, J. Soviet Math. 9(1978) 697–749.
- [25] P. L. Lion, Mathematics Topics in Fluid Mechanics, in: Oxford Lecture Series in Mathematics and its applications, vol. 1, The Clarendon Press University Press, New York, 1996.
- [26] A. Meyer, Wavelets, Paraproducts and Navier-Stokes equations, Current developments in mathematics, International press, 1996.
- [27] A. Matsumura and T. Nishita, The initial value problem for the equations of motion of viscous and heat conductive gses, J. Math. Kyoto Univ. 20(1)(1980) 67–104.
- [28] J. Nash, Le problème de cauchy pour les équations différentielles d’un fluide général, Bulletin de la Société Mathématique de France 90(1962) 487–497.
- [29] A. I. Volpert, S. I. Hudjaev, On the cauchy probelm for composite systems of nonlinear differential equations, Math. USSR-Sb 16(1972) 517–544.
- [30] Z. P. Xin, Blow up of smooth solutions to the compressible Navier-Stokes equation with compact density, Comm. Pure Appl. Math. 51(1998) 229–240.

THE GRADUATE SCHOOL OF CHINA ACADEMY OF ENGINEERING PHYSICS, P. O. BOX 2101,
BEIJING 100088, PR CHINA,

E-mail address: bian_dongfen@mail.com

INSTITUTE OF APPLIED PHYSICS AND COMPUTATIONAL MATHEMATICS, P. O. BOX 8009, BEIJING
100088, PR CHINA.

E-mail address: gbl@iapcm.ac.cn